

# Aldo-keto reductase (AKR) superfamily: Genomics and annotation

Rebekka D. Mindnich and Trevor M. Penning\*

Center for Excellence in Environmental Toxicology, Department of Pharmacology, University of Pennsylvania, School of Medicine, Philadelphia, PA 19104-6084, USA

\*Correspondence to: Tel: +1 215 898 9445; Fax: +1 215 573 2236; E-mail: penning@mail.med.upenn.edu

Date received (in revised form): 22nd May 2009

## Abstract

Aldo-keto reductases (AKRs) are phase I metabolising enzymes that catalyse the reduced nicotinamide adenine dinucleotide (phosphate) (NAD(P)H)-dependent reduction of carbonyl groups to yield primary and secondary alcohols on a wide range of substrates, including aliphatic and aromatic aldehydes and ketones, ketoprostaglandins, ketosteroids and xenobiotics. In so doing they functionalise the carbonyl group for conjugation (phase II enzyme reactions). Although functionally diverse, AKRs form a protein superfamily based on their high sequence identity and common protein fold, the  $(\alpha/\beta)_8$ -barrel structure. Well over 150 AKR enzymes, from diverse organisms, have been annotated so far and given systematic names according to a nomenclature that is based on multiple protein sequence alignment and degree of identity. Annotation of non-vertebrate AKRs at the National Center for Biotechnology Information or Vertebrate Genome Annotation (vega) database does not often include the systematic nomenclature name, so the most comprehensive overview of all annotated AKRs is found on the AKR website (<http://www.med.upenn.edu/akr/>). This site also hosts links to more detailed and specialised information (eg on crystal structures, gene expression and single nucleotide polymorphisms [SNPs]). The protein-based AKR nomenclature allows unambiguous identification of a given enzyme but does not reflect the wealth of genomic and transcriptomic variation that exists in the various databases. In this context, identification of putative new AKRs and their distinction from pseudogenes are challenging. This review provides a short summary of the characteristic features of AKR biochemistry and structure that have been reviewed in great detail elsewhere, and focuses mainly on nomenclature and database entries of human AKRs that so far have not been subject to systematic annotation. Recent developments in the annotation of SNP and transcript variance in AKRs are also summarised.

**Keywords:** carbonyl reduction, nomenclature, pseudogene, SNP, splice variant

## Introduction

Aldo-keto reductases (AKRs) form a superfamily of proteins characterised by their common three-dimensional structure and reaction mechanism in catalysing the reduced nicotinamide adenine dinucleotide (phosphate) (NAD(P)H)-dependent oxido-reduction of carbonyl groups.<sup>1,2</sup> More than 150 enzymes have been identified in a wide range of organisms, including prokaryotes and eukaryotes such as plants, fungi and vertebrates. Based on

sequence identity, the proteins fall into 15 different families, termed AKR1–AKR15, and some families (eg AKR1) contain multiple subfamilies.<sup>1,3,4</sup> Fourteen human AKRs (Table 1), which fall into families 1, 6 and 7, have been annotated and their specific structural and kinetic features, as well as substrate and genomic characteristics, have been recently and comprehensively reviewed.<sup>5,6</sup>

AKRs are, for the most part, cytosolic and are active as monomers. So far, only AKR2, AKR6

**Table 1.** Systematic and trivial names of human Aldo-keto reductases (AKRs)

Systematic name	Trivial name/often-used abbreviations
AKR1A1	Aldehyde reductase; ALR; ALDR1
AKR1B1	Aldose reductase; AR, ALR2
AKR1B10	Small intestine-like aldose reductase; ARL-1
AKR1C1	Dihydrodiol dehydrogenase 1; 20 $\alpha$ , (3 $\alpha$ )-Hydroxysteroid dehydrogenase
AKR1C2	Dihydrodiol dehydrogenase 2; 3 $\alpha$ -Hydroxysteroid dehydrogenase, type III
AKR1C3	3 $\alpha$ -Hydroxysteroid dehydrogenase, type II
AKR1C4	Dihydrodiol dehydrogenase 4; chlordecone reductase; 3 $\alpha$ -hydroxysteroid dehydrogenase, type I
AKR1D1	$\Delta^4$ -3-Ketosteroid-5 $\beta$ -reductase
AKR1E2	Aldo-keto reductase family I, member C-like 2; AKR1CL2
AKR6A3	KCNAB 1; hKvbeta3; potassium voltage-gated channel, shaker-related subfamily, $\beta$ - member 1
AKR6A5	KCNAB 2; hKvbeta2; potassium voltage-gated channel, shaker-related subfamily, $\beta$ - member 2
AKR6A9	KCNAB 3; potassium voltage-gated channel, shaker-related subfamily, $\beta$ - member 3
AKR7A2	Aflatoxin aldehyde reductase; AFAR1
AKR7A3	Aflatoxin aldehyde reductase; AFAR2

and AKR7 family members show evidence of forming multimers, although it is unclear whether quaternary structure is essential for enzyme activity.<sup>7–9</sup> The single amino acid chain of, on average, 320 residues folds into a  $(\alpha/\beta)_8$ -barrel, an evolutionarily highly conserved structure originally observed in triose-phosphate isomerase (TIM) and therefore also known as a TIM barrel structure. In contrast to another 32 superfamilies adopting this motif, AKRs do not require metal ions for functionality, and bind NAD(P)H cofactor in an

extended *anti*-conformation so that they can catalyse 4-*pro-R* hydride transfer and thus belong to the large group of A-face oxidoreductases.

The spectrum of AKR substrates is wide and comprises aldehydes, ketones, monosaccharides, ketosteroids, ketoprostaglandins, bile acid precursors, chemical carcinogens and their metabolites — for example, aflatoxin dialdehydes, nicotine-derived 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and *trans*-dihydrodiols of polycyclic aromatic hydrocarbons. Substrate recognition and binding takes place at the C-terminal end of the ellipsoid  $(\alpha/\beta)_8$ -barrel body through interaction with amino acids within the three highly variable loops protruding out of the barrel structure.<sup>2,10,11</sup> Indeed, sequence conservation among AKRs is lowest in these loops, compared with the  $\alpha$ -helices and  $\beta$ -strands forming the barrel structure itself, and suggests that AKRs evolved from a common ancestor by duplication and subsequent divergence to accommodate the binding of the structurally diverse substrates.

Duplication and diversification indeed seem to have been the driving force in the formation of vertebrate AKRs. AKR1B, AKR1C and AKR7A family members appear to be vertebrate specific and in the human genome their genes cluster on chromosomes 7, 10 and 1, respectively.<sup>5</sup> Furthermore, the potential of these three subfamilies to give rise to new functional AKRs is reflected in the existence of several pseudogenes and putative new isoforms in the human genome. By contrast, the genes of all three human AKR6 family members reside in different chromosomal locations, show a high degree of conservation, even to plant AKR6 isoforms, and do not give rise to functional or non-functional (pseudogene) gene duplications.

## Biochemistry of AKRs

Almost all AKRs catalyse the reduction of aldehydes and ketones to primary and secondary alcohols, respectively, by use of a reduced nicotinamide cofactor. The reactions involve an ordered bi-bi kinetic mechanism and general acid-base catalysis.<sup>12–14</sup> The cofactor binds first, followed by

tethering of the substrate. In the reduction direction, the 4-*pro-R* hydride from NAD(P)H is transferred to the substrate carbonyl group, followed by protonation of the carbonyl oxygen by a conserved tyrosine acting as a general acid.<sup>15–17</sup> Oxidation proceeds in the reverse sequence with the tyrosine acting as a general base.


Asp 50, Tyr 55, Lys 84 and His 117 form the catalytic tetrad (based on residue numbering in AKR1C9),<sup>2,18,19</sup> where the Tyr is the general acid-base.<sup>17,20</sup> Conservation of these residues is strong and underlines their role in catalysis. Asp 50 is present in 99 per cent, Tyr 55 in 97 per cent, Lys 84 in 97 per cent and His 117 in 88 per cent of all annotated AKRs on the AKR website. Consistent substitution of these key amino acids, however, especially of the histidine, is a characteristic for some AKR subfamilies. In AKR1D isoforms, which catalyse the 5 $\beta$ -reduction of  $\Delta^4$ -3-ketosteroids, glutamate replaces His 117.<sup>2,21</sup> Interestingly, mutational analysis in AKR1C9 demonstrates that this single amino acid substitution is sufficient to convert this 3-ketosteroid reductase to a steroid 5 $\beta$  double-bond reductase.<sup>22,23</sup> The reverse change of function has been hypothesised for the respective (E120H) mutation in AKR1D1.<sup>24</sup> In all AKR6A isoforms, the conserved histidine is changed to asparagine. Based on crystal structure comparisons, Di Costanzo *et al.* hypothesised that in AKR6A isoforms the amino acid in this position is involved in substrate discrimination.<sup>25</sup> AKR6A isoforms, better known as voltage-gated potassium channel  $\beta$ -subunits (Kv $\beta$  or KCNAB), accept a wide range of substrates, but their catalytic efficiency is significantly lower than other AKRs.<sup>26,27</sup> It has been suggested that they may primarily function as redox state sensors in the regulation of potassium channel activity.<sup>27–29</sup> The replacement of histidine with asparagine may provide more space to allow for the optimal positioning of bulky lipid and phospholipid aldehydes, formed as products of lipid peroxidation and oxidative stress.

Although these four conserved amino acids seem to play a crucial role in the catalytic mechanism, targeted mutation studies (eg in AKR1B1<sup>15</sup> and AKR1C9<sup>30</sup>) did not indicate a complete loss in enzymatic activity and suggested that, in some

instances, proximity of the hydride donor to the acceptor is sufficient to catalyse a reaction. Furthermore, some AKRs, such as  $\rho$ -crystallins in amphibian lenses (AKR1C10), appear to have been employed in this tissue basically for structural reasons, having lost almost all enzymatic activity despite unaltered characteristics of the cofactor and substrate binding site.<sup>31</sup>

## Nomenclature

The nomenclature for the AKR superfamily, introduced and accepted by the Hugo Genome Nomenclature Committee (HGNC) in 1997, builds solely on sequence and not on functional similarity. Furthermore, systematic annotation through the AKR website requires proof of functionality of the protein in question. This evidence may be RNA or enzyme based, and aims to avoid the annotation of non-functional pseudogenes or predicted genes of unknown physiological function. Sequence submission for the assignment of a systematic AKR superfamily name should be sent directly to the AKR website (<http://www.med.upenn.edu/akr/>).<sup>3,4</sup> The basic principle in naming a new AKR follows that for the cytochrome P (CYP) superfamily.<sup>32,33</sup> All members of the superfamily are annotated with the suffix AKR. The 15 families (Figure 1) into which annotated AKRs fall have less than 40 per cent amino acid identity with each other. Subfamilies, denoted by a capital roman letter, share up to 60 per cent sequence identity. Individual members in the subfamily are numbered according to the chronology of sequence submission to the AKR website. Thus, AKR1A1 (human aldehyde reductase) belongs to family 1, subfamily A and is the first unique protein in the subfamily. If two sequences share 97 per cent or more sequence identity, they are considered as alleles and marked by an additional small roman letter (eg AKR1C10a and AKR1C10b). Two sequences sharing this high degree of identity are not regarded as alleles if they reside in different genomic locations, have distinct 3' untranslated regions or differ in their enzymatic activities. It was also recommended that the corresponding gene be



given the same name as the protein, but written in italics.

The nomenclature has the advantage that it assigns a unique gene name to each single AKR through establishing its amino acid sequence similarity to other families and subfamilies. In this way, it is independent of enzyme properties, species of origin and functional homologies. Therefore, it is not surprising that most families at the same time constitute clusters of closely related organisms rather than similar function, and that an AKR of a more distantly related organism will fall into a different subfamily (Figure 1). For example, bacterial AKRs form their own families with multiple subfamilies owing to the high diversification that has occurred in the 3,900 million years since bacteria and archaea split. At the same time, vertebrate, but especially mammalian, AKR sequences (mammals appeared about 300 million years ago) form sequence clusters with multiple, highly similar

members in each subfamily. Inspection of the vertebrate AKR sequences reveals that the AKR1 family evolved separately from families AKR6 and AKR7. Furthermore, only potassium channel  $\beta$ -subunit sequences have been so well conserved during evolution that plant and arthropod sequences both cluster in the AKR6 family.

## A search for AKRs on NCBI, HGNC and Ensembl

Although the new nomenclature came into effect nearly 12 years ago—and has been widely accepted by the AKR research community—it has not yet become widely adopted by database annotators. This is especially true for non-mammalian AKRs. Of the 160 proteins annotated on the AKR website (<http://www.med.upenn.edu/akr/>), 62 give problematic results if searched with their systematic name against the National Center for Biotechnology Information (NCBI) database. For 45 AKRs, not a single database entry can be found. This illustrates that regular updates of protein and nucleotide annotations have failed to notice and acknowledge the proposed systematic nomenclature.

Annotation and nomenclature of human AKRs, however, are far more consistent. A database search for human AKRs at the National Institutes of Health (<http://www.ncbi.nlm.nih.gov/>), the European Molecular Biology Laboratory ([www.embl.org](http://www.embl.org/)), the Sanger Institute (<http://vega.sanger.ac.uk/index.html>) and the HGNC (<http://www.genenames.org/>) identified 13 of the 14 proteins listed on the AKR website (Table 1). AKR1E2, which appears in all four databases under its synonym AKR1CL2, was originally annotated as LoopADR in 2000 and later named AKR1C like 2 (AKR1CL2), based on its sequence similarity to AKR1C isoforms. In 2003 and 2004, two publications described the enzymatic activity of this protein towards 9,10-phenanthrenequinone, but not steroid hormones, and termed it human testis-specific AKR (htAKR), based on its restricted expression pattern.<sup>34,35</sup> The protein shares 74 per cent sequence identity with the previously described

murine AKR1E1<sup>36</sup> and hence was systematically named AKR1E2.

## Unclassified and predicted human AKR database entries

The database search for human AKRs at NCBI also revealed a number of other sequences annotated in context with AKRs. The total of 58 database entries comprised the 14 systematically annotated and well-described human AKRs and 12 separate annotations of pseudogenes. The search also revealed that in 22 cases, the respective protein belonged to a different protein family but appears in a biological context with an AKR. The remaining ten database entries (Table 2) resemble

unclassified or predicted human AKR sequences that are related to three different AKR subfamilies: AKR1C, AKR1B and AKR7A.

*AKR1CL1* is located between *AKR1C3* and *AKR1C4* on chromosome 10. Ambiguous reports at NCBI describe two different forms of the gene. In one case, exons 1–4 of eight exons encode a 129-amino acid peptide (Accession No. NP001007537.1). This protein includes the complete catalytic tetrad but appears to be too short to be a functional AKR. In the second case, *AKR1CL1*, similarly to the other four *AKR1C* members, consists of nine exons and encodes a complete 326-amino acid protein (Accession No. EAW86451) with 68 per cent sequence identity to *AKR1C1-4*. The shorter version exactly matches

**Table 2.** Non-classified human AKRs in the NCBI database

Name	Alternative names	Gene ID <sup>a</sup>	Chromosomal location	Related to	Probable identity
<i>AKR1B11</i>	<i>ARL, HSI</i>	9405	7	AKR1B	Variance of <i>AKR1B10</i>
<i>AKR1B10L</i>	<i>tcag7.1260</i>	441282	7q33	AKR1B	New AKR or pseudogene of <i>AKR1B10</i>
<i>LOC340888</i>		340888	10q21.3	AKR1B	Retro-transposed <i>AKR1B10</i> pseudogene
<i>LOC643582</i>		643582	18q22.1	AKR1B	Retro-transposed <i>AKR1B10</i> pseudogene
<i>AKR1CL1</i>	<i>RAKc</i>	340811	10p15.1	AKR1C	New <i>AKR1C</i> or pseudogene
<i>AKR1C-pseudo</i>		266745	10p15–10p14	AKR1C	Variance of <i>AKR1C2</i>
<i>LOC648517</i>		648517	Unclear	AKR1C	<i>AKR1C</i> pseudogene or transcript variant
<i>LOC100134257</i>		100134257	Unclear	AKR1C	<i>AKR1C</i> pseudogene or transcript variant
<i>LOC643789</i>		643789	10p15.1	AKR1C	Pseudogene
<i>LOC648947</i>		648947	10p15.1	AKR1C	<i>AKR1CL1</i> pseudogene
<i>tAKR</i>		389932	10p15.1	AKR1C	<i>AKR1C</i> pseudogene
<i>AKR7L</i>	<i>AKR7A4, AFAR3</i>	246181	1p35–1p36.1	AKR7A	New <i>AKR7A</i> or pseudogene
<i>AFARPI</i>		246182	1p12	AKR7A	Retro-transposed <i>AKR7</i> pseudogene

<sup>a</sup>At NCBI (<http://www.ncbi.nlm.nih.gov>).

residues 1–129 of the longer form and resembles a differently transcribed and processed version of the same gene. A few messenger RNA (mRNA) sequences are annotated that match *AKR1CL1* with deletions and frameshifts, but a Blast search with the longer version against the expressed sequence tag (EST) database does not yield any *AKR1CL1* transcripts. Therefore, it remains unclear whether *AKR1CL1* is a processed pseudogene or a new functional AKR1C member.

*AKR1C-pseudo* is an old, discontinued entry that resembles *AKR1C2*. *AKR1C2* shares 97 per cent sequence identity with *AKR1C1* but exhibits a unique expression pattern and substrate specificity.<sup>37–41</sup> *LOC648517* and *LOC100134257* contain sequence elements that completely match exons of *AKR1C1* and *AKR1C2*. Their chromosomal location could not be determined, however, and therefore it is not clear whether they might be transcript variants of either *AKR1C1* or *AKR1C2*, or are separate pseudogenes.

Four entries are related to *AKR1B10*. While *AKR1B11* is a discontinued record replaced by *AKR1B10*, *LOC340888* and *LOC643582* are Gnomon predictions and appear to be retro-transposed pseudogenes of *AKR1B10*. *AKR1B10L* resides next to *AKR1B10* on chromosome 7 and also consists of ten exons. The two proteins share 91 per cent amino acid identity. The presence of a single EST from placenta (BX350113.2) that completely matches residues 23–209 of the predicted 316-amino acid *AKR1B10L* transcript suggests that *AKR1B10L* could indeed resemble a new and thus far uncharacterised human AKR family member. Alternatively, the presence of the mRNA could result from a processed pseudogene.

Two additional database entries relate to the AKR7 family. *AKR7L* clusters together with *AKR7A2* and *AKR7A3* on chromosome 1p35–1p36.1.<sup>3,42</sup> Alternative transcription leads to two different transcript variants. Variant 1 is the longer and comprises seven exons that encode a 331-amino acid protein. This form may indeed resemble a new functional AKR that has 92 per cent and 88 per cent identity with *AKR7A3* and *AKR7A2*, respectively. Variant 2 lacks exons 4 and 5

and codes for a probably non-functional alternative peptide. This deletion results in a frameshift-eliminating sequence similarity to AKRs in the last 80 amino acids of the 250-amino acid peptide. The *AKR7* family pseudogene (*AFARP1*) is a retro-transposed *AKR7*, where the complete processed mRNA has reintegrated into the genome.

In addition, Barski *et al.*<sup>5</sup> report on three putative AKRs not identified in the above-described database search. The database entry *LOC643789* was originally designated as the *AKR1C* pseudogene but was discontinued in March 2008. *LOC648947* is a predicted gene with nine exons, encodes a 316-amino acid protein and shares 82 per cent sequence identity with the longer version of *AKR1CL1*; however, this protein sequence harbours a stop codon following amino acid 199. At least three mRNA sequences for this gene are present in the EST database. They do not completely match the predicted protein sequence of *LOC648947*, and a region corresponding to amino acids 192–227 is deleted. As this deletion covers a region conserved in all *AKR1C* isoforms, it is very likely that *LOC648947* represents a processed pseudogene of *AKR1CL1*. Finally, *tAKR* shows similarity to other *AKR1C* isoforms and the gene is located between *AKR1CL2* and *AKR1C1* on chromosome 10. Original annotation predicts a gene of 11 exons, with exons 3–11 encoding a putative protein with more than 300 amino acids. A transcript exists only for the first 125 amino acids, however, and hence *tAKR* is thus far classified as a processed pseudogene.

In summary, systematic integration of these AKR database entries into the AKR nomenclature is still pending owing to a lack of sufficient experimental data to support an assignment.

## Annotated variants of AKRs

Efforts have been made systematically to describe genetic variants and splice isoforms in annotated human AKRs and to identify the location and classification of putative pseudogenes (Table 3). Data on single nucleotide polymorphisms (SNPs) are accumulating rapidly, and the SNP database

**Table 3.** Genomic and transcriptomic variations in annotated human AKRs

AKR	Gene ID <sup>a</sup>	Chromosomal location	No. of SNPs <sup>b</sup> (in gene / coding region / non-synonymous / frameshift)	SNP frequency (per 1 kilobase: in gene / in coding region)	Peptide variants	Additional processed transcripts	Pseudogenes on vega (HGNC)
<i>IA1</i>	10327	1p33–p32	154 / 9 / 5 / 1	8.01 / 9.2	3	10	0
<i>IB1</i>	231	7q35	160 / 13 / 9 / 0	9.53 / 13.67	3	12	5 (8)
<i>IB10</i>	57016	7q33	127 / 16 / 5 / 0	9.65 / 16.82	1	3	4
<i>IC1</i>	1645	10p15–p14	160 / 16 / 6 / 0	10.88 / 16.46	3	2	0
<i>IC2</i>	1646	10p15–p14	353 / 16 / 8 / 0	12.5 / 16.46	1	1	0
<i>IC3</i>	8644	10p15–p14	261 / 23 / 13 / 2	19.61 / 23.66	1	3	0
<i>IC4</i>	1109	10p15–p14	198 / 7 / 6 / 0	8.95 / 7.2	1	1	0
<i>ID1</i>	6718	7q32–q33	275 / 1 / 0 / 0	6.57 / 1.02	4	2	2 (1)
<i>IE2</i>	83592	10p15.1	211 / 12 / 7 / 0	9.66 / 12.46	2	4	0
<i>6A3</i>	7881	3q26.1	1888 / 1 / 0 / 0	4.51 / 0.81	10	0	0
<i>6A5</i>	8514	1p36.3	414 / 8 / 2 / 1	7.15 / 7.25	8	6	0
<i>6A9</i>	9196	17p13.1	42 / 3 / 1 / 2	6.24 / 2.47	1	2	0
<i>7A2</i>	8574	1p36.13	71 / 10 / 8 / 0	8.68 / 9.26	1	0	4
<i>7A3</i>	22977	1p36.13	67 / 12 / 5 / 0	10.76 / 12.05	1	0	0

<sup>a</sup>At NCBI (<http://www.ncbi.nlm.nih.gov>).

<sup>b</sup>Deduced from the vega database (<http://vega.sanger.ac.uk/index.html>).

The table does not include the putative transcript variants and pseudogenes listed in Table 2.

build 130 (NCBI), as of 30th April 2009, contains over 17.8 million SNP clusters, with over 7.3 million SNPs located in genes. Nomenclature for SNPs in *AKRs* is based on that for sulfotransferase (*SULT*) genes<sup>43</sup> and takes only coding SNPs into account.<sup>4</sup> Systematic annotation of the SNP is marked by the gene name followed by an asterisk, an Arabic number indicating synonymous ('1') or non-synonymous ('2'–'n') character and a letter distinguishing between suballeles; the nomenclature is chronologically based. While this seems to be a feasible approach for synonymous and non-synonymous SNPs, it neglects SNPs leading to frameshifts and those in the non-coding gene region that have the potential to alter the transcription efficiency, splicing and stability of mRNA. For all *AKRs* SNP frequency is significantly higher than the estimated 1 SNP/kilobase.<sup>44,45</sup> Although the total number of SNPs per gene is only a snapshot

in this time of extremely rapid data growth, it appears that some *AKRs* (eg *AKR1D1*, *AKR6A3* and *AKR6A9*) have accumulated significantly fewer single nucleotide variations in their coding region than others.

In addition to SNPs, other variations have been annotated over the years. The most well known are pseudogenes, and one or several have been identified for a few *AKRs*. For example, the pseudogene forms of *AKR1B1* are well documented and systematically annotated as *AKR1B1P1-8*, where P stands for pseudogene. For example, *AKR1B1P4* denotes the fourth pseudogene annotated for *AKR1B1*. Analysis of the non-systematically named *AKRs* (Table 2) suggests that many more pseudogenes, in addition to the few already annotated, might exist. This may also be the case for alternative splice forms, where the vega database (<http://vega.sanger.ac.uk/index.html>) from the Sanger Institute offers the most systematic

annotation approach. Gene transcripts are numbered in the order of their annotation and designated with the suffix '-n' added to the specific gene name. For example, *AKR1C1* has five different transcripts, annotated as *AKR1C1-001* to *AKR1C1-005*. For half of the human *AKRs*, transcript processing leads to one or more alternative peptides. Furthermore, non-coding RNAs have been identified for 11 of the 14 genes. While, in the past, these fragments have been ignored, being perceived as resulting from errors in transcription or splicing, increasing evidence suggests that they play important regulatory functions (eg in gene expression, imprinting, RNA stability, splicing and translation).<sup>46</sup> Their role in *AKR* expression and function remains to be elucidated.

## Summary

Proteins of the *AKR* superfamily are present in all phyla and are characterised by their common three-dimensional structure and reaction mechanism. Fourteen human *AKRs* have so far been annotated with systematic gene names in accordance with a systematic nomenclature provided through the *AKR* website that is based on amino acid sequence similarity and proof of protein functionality. Additional non-systematic annotations of human *AKRs* are present in the NCBI database that may resemble new, uncharacterised proteins or pseudogenes of existing *AKRs*. Furthermore, the rapid and continuous increase in data on SNPs and transcript variants challenges a systematic annotation of these phenomena and adds to the complexity of defining the functionality of a given gene product. Despite these complexities, the *AKR* website provides the most comprehensive and up-to-date portal for tracking the development of *AKR* annotation and provides links to in-depth information on each single *AKR* superfamily member in other databases.

## Acknowledgments

The writing of this paper was funded in part by NIH grants P30-ES013508, R01-DK47015, R01-CA90744 and R01-CA39504 awarded to T.M.P.

## References

1. Jez, J.M., Flynn, T.G. and Penning, T.M. (1997), 'A nomenclature system for the aldo-keto reductase superfamily', *Adv. Exp. Med. Biol.* Vol. 414, pp. 579–600.
2. Jez, J.M., Bennett, M.J., Schlegel, B.P., Lewis, M. and Penning, T.M. (1997), 'Comparative anatomy of the aldo-keto reductase superfamily', *Biochem. J.* Vol. 326, pp. 625–636.
3. Hyndman, D., Bauman, D.R., Heredia, V.V. and Penning, T.M. (2003), 'The aldo-keto reductase superfamily homepage', *Chem. Biol. Interact.* Vol. 143–144, pp. 621–631.
4. Drury, J.E., Hyndman, D., Jin, Y. *et al.* (2007), 'The aldo-keto reductase superfamily homepage: 2006 update', In: Weiner, H., Maser, E., Lindhal, R. *et al.* (eds), *Enzymology and Molecular Biology of Carbonyl Metabolism*, Purdue Press, West Lafayette, IN, pp. 184–197.
5. Barski, O.A., Tipparaju, S.M. and Bhatnagar, A. (2008), 'The aldo-keto reductase superfamily and its role in drug metabolism and detoxification', *Drug Metab. Rev.* Vol. 40, pp. 553–624.
6. Jin, Y. and Penning, T.M. (2007), 'Aldo-keto reductases and bioactivation/detoxification', *Annu. Rev. Pharmacol. Toxicol.* Vol. 47, pp. 263–292.
7. Kozma, E., Brown, E., Ellis, E.M. and Laphorn, A.J. (2002), 'The crystal structure of rat liver AKR7A1. A dimeric member of the aldo-keto reductase superfamily', *J. Biol. Chem.* Vol. 277, pp. 16285–16293.
8. Kavanagh, K.L., Klimacek, M., Nidetzky, B. and Wilson, D.K. (2002), 'The structure of apo and holo forms of xylose reductase, a dimeric aldo-keto reductase from *Candida tenuis*', *Biochemistry* Vol. 41, pp. 8785–8795.
9. Gulbis, J.M., Mann, S. and Mackinnon, R. (1999), 'Structure of a voltage-dependent K<sup>+</sup> channel  $\beta$  subunit', *Cell* Vol. 97, pp. 943–952.
10. Bennett, M.J., Albert, R.H., Jez, J.M., Ma, H. *et al.* (1997), 'Steroid recognition and regulation of hormone action: Crystal structure of testosterone and NADP<sup>+</sup> bound to 3 $\alpha$ -hydroxysteroid/dihydrodiol dehydrogenase', *Structure* Vol. 5, pp. 799–812.
11. Couture, J.F., De Jesus-Tran, K.P., Roy, A.M., Cantin, L. *et al.* (2005), 'Comparison of crystal structures of human type 3 3 $\alpha$ -hydroxysteroid dehydrogenase reveals an "induced-fit" mechanism and a conserved basic motif involved in the binding of androgen', *Protein Sci.* Vol. 14, pp. 1485–1497.
12. Askonas, L.J., Ricigliano, J.W. and Penning, T.M. (1991), 'The kinetic mechanism catalysed by homogeneous rat liver 3 $\alpha$ -hydroxysteroid dehydrogenase. Evidence for binary and ternary dead-end complexes containing non-steroidal anti-inflammatory drugs', *Biochem. J.* Vol. 278, pp. 835–841.
13. Neuhauser, W., Haltrich, D., Kulbe, K.D. and Nidetzky, B. (1997), 'NAD(P)H-dependent aldose reductase from the xylose-assimilating yeast *Candida tenuis*. Isolation, characterization and biochemical properties of the enzyme', *Biochem. J.* Vol. 326, pp. 683–692.
14. Trauger, J.W., Jiang, A., Stearns, B.A. and Logrosso, P.V. (2002), 'Kinetics of allopregnanolone formation catalyzed by human 3 $\alpha$ -hydroxysteroid dehydrogenase type III (AKR1C2)', *Biochemistry* Vol. 41, pp. 13451–13459.
15. Bohren, K.M., Grimshaw, C.E., Lai, C.J., Harrison, D.H. *et al.* (1994), 'Tyrosine-48 is the proton donor and histidine-110 directs substrate stereochemical selectivity in the reduction reaction of human aldose reductase: Enzyme kinetics and crystal structure of the Y48H mutant enzyme', *Biochemistry*, Vol. 33, pp. 2021–2032.
16. Penning, T.M. (1999), 'Molecular determinants of steroid recognition and catalysis in aldo-keto reductases. Lessons from 3 $\alpha$ -hydroxysteroid dehydrogenase', *J. Steroid Biochem. Mol. Biol.* Vol. 69, pp. 211–225.
17. Schlegel, B.P., Jez, J.M. and Penning, T.M. (1998), 'Mutagenesis of 3 $\alpha$ -hydroxysteroid dehydrogenase reveals a "push-pull" mechanism for proton transfer in aldo-keto reductases', *Biochemistry* Vol. 37, pp. 3538–3548.
18. Jez, J.M., Flynn, T.G. and Penning, T.M. (1997), 'A new nomenclature for the aldo-keto reductase superfamily', *Biochem. Pharmacol.* Vol. 54, pp. 639–647.
19. Bennett, M.J., Schlegel, B.P., Jez, J.M., Penning, T.M. and Lewis, M. (1996), 'Structure of 3 $\alpha$ -hydroxysteroid/dihydrodiol dehydrogenase complexed with NADP<sup>+</sup>', *Biochemistry* Vol. 35, pp. 10702–10711.



20. Grimshaw, C.E., Bohren, K.M., Lai, C.J. and Gabbay, K.H. (1995), 'Human aldose reductase: pK of tyrosine 48 reveals the preferred ionization state for catalysis and inhibition', *Biochemistry* Vol. 34, pp. 14374–14384.
21. Kondo, K.H., Kai, M.H., Setoguchi, Y., Eggertsen, G. *et al.* (1994), 'Cloning and expression of cDNA of human  $\Delta^4$ -3-oxosteroid 5 $\beta$ -reductase and substrate specificity of the expressed enzyme', *Eur. J. Biochem.* Vol. 219, pp. 357–363.
22. Penning, T.M., Ma, H. and Jez, J.M. (2001), 'Engineering steroid hormone specificity into aldo-keto reductases', *Chem. Biol. Interact.* Vol. 130–132, pp. 659–671.
23. Jez, J.M. and Penning, T.M. (1998), 'Engineering steroid 5 $\beta$ -reductase activity into rat liver 3 $\alpha$ -hydroxysteroid dehydrogenase', *Biochemistry* Vol. 37, pp. 9695–9703.
24. Drury, J.E. and Penning, T.M. (2007), ' $\Delta^4$ -3-ketosteroid 5 $\beta$ -reductase (AKR1D1): Properties and role in bile acid synthesis', In: Weiner, H., Maser, E., Lindhal, R. *et al.* (eds), *Enzymology and Molecular Biology of Carbonyl Metabolism*, Purdue Press, West Lafayette, IN, pp. 332–340.
25. Di Costanzo, L., Penning, T.M. and Christianson, D.W. (2009), 'Aldo-keto reductases in which the conserved catalytic histidine is substituted', *Chem. Biol. Interact.* Vol. 178, pp. 127–133.
26. Tipparaju, S.M., Barski, O.A., Srivastava, S. and Bhatnagar, A. (2008), 'Catalytic mechanism and substrate specificity of the  $\beta$ -subunit of the voltage-gated potassium channel', *Biochemistry* Vol. 47, pp. 8840–8854.
27. Weng, J., Cao, Y., Moss, N. and Zhou, M. (2006), 'Modulation of voltage-dependent Shaker family potassium channels by an aldo-keto reductase', *J. Biol. Chem.*, Vol. 281, pp. 15194–15200.
28. Tipparaju, S.M., Saxena, N., Liu, S.Q., Kumar, R. and Bhatnagar, A. (2005), 'Differential regulation of voltage-gated K<sup>+</sup> channels by oxidized and reduced pyridine nucleotide coenzymes', *Am. J. Physiol. Cell Physiol.* Vol. 288, pp. C366–C376.
29. Pan, Y., Weng, J., Cao, Y., Bhosle, R.C. and Zhou, M. (2008), 'Functional coupling between the Kv1.1 channel and aldoketoreductase Kv $\beta$ 1', *J. Biol. Chem.*, Vol. 283, pp. 8634–8642.
30. Schlegel, B.P., Ratnam, K. and Penning, T.M. (1998), 'Retention of NADPH-linked quinone reductase activity in an aldo-keto reductase following mutation of the catalytic tyrosine', *Biochemistry* Vol. 37, pp. 11003–11011.
31. Fujii, Y., Watanabe, K., Hayashi, H., Urade, Y. *et al.* (1990), 'Purification and characterization of rho-crystallin from Japanese common bullfrog lens', *J. Biol. Chem.* Vol. 265, pp. 9914–9923.
32. Dayhoff, M.O., Barker, W.C. and Hunt, L.T. (1983), 'Establishing homologies in protein sequences', *Methods Enzymol.* Vol. 91, pp. 524–545.
33. Nebert, D.W., Adesnik, M., Coon, M.J. *et al.* (1987), 'The P450 gene superfamily: recommended nomenclature', *DNA* Vol. 6, No. 1, pp. 1–11.
34. Nishinaka, T., Azuma, Y., Ushijima, S., Miki, S. *et al.* (2003), 'Human testis specific protein: A new member of aldo-keto reductase superfamily', *Chem. Biol. Interact.* Vol. 143–144, pp. 299–305.
35. Azuma, Y., Nishinaka, T., Ushijima, S., Soh, J. *et al.* (2004), 'Characterization of htAKR, a novel gene product in the aldo-keto reductase family specifically expressed in human testis', *Mol. Hum. Reprod.* Vol. 10, pp. 527–533.
36. Bohren, K.M., Barski, O.A. and Gabbay, K.H. (1997), 'Characterization of a novel murine aldo-keto reductase', *Adv. Exp. Med. Biol.* Vol. 414, pp. 455–464.
37. Ciaccio, P.J. and Tew, K.D. (1994), 'cDNA and deduced amino acid sequences of a human colon dihydrodiol dehydrogenase', *Biochim. Biophys. Acta* Vol. 1186, pp. 129–132.
38. Qin, K.N., New, M.I. and Cheng, K.C. (1993), 'Molecular cloning of multiple cDNAs encoding human enzymes structurally related to 3 $\alpha$ -hydroxysteroid dehydrogenase', *J. Steroid Biochem. Mol. Biol.* Vol. 46, pp. 673–679.
39. Stolz, A., Hammond, L., Lou, H., Takikawa, H. *et al.* (1993), 'cDNA cloning and expression of the human hepatic bile acid-binding protein. A member of the monomeric reductase gene family', *J. Biol. Chem.* Vol. 268, pp. 10448–10457.
40. Penning, T.M., Jin, Y., Heredia, V.V. and Lewis, M. (2003), 'Structure-function relationships in 3 $\alpha$ -hydroxysteroid dehydrogenases: A comparison of the rat and human isoforms', *J. Steroid Biochem. Mol. Biol.* Vol. 85, pp. 247–255.
41. Penning, T.M., Jin, Y., Steckelbroeck, S., Lanisnik-Rizner, T. and Lewis, M. (2004), 'Structure-function of human 3 $\alpha$ -hydroxysteroid dehydrogenases: Genes and proteins', *Mol. Cell Endocrinol.* Vol. 215, pp. 63–72.
42. Praml, C., Savelyeva, L. and Schwab, M. (2003), 'Aflatoxin B1 aldehyde reductase (AFAR) genes cluster at 1p35-1p36.1 in a region frequently altered in human tumour cells', *Oncogene* Vol. 22, pp. 4765–4773.
43. Blanchard, R.L., Freimuth, R.R., Buck, J., Weinsilbourn, R.M. and Coughtrie, M.W. (2004), 'A proposed nomenclature system for the cytosolic sulfotransferase (SULT) superfamily', *Pharmacogenetics* Vol. 14, pp. 199–211.
44. Kwok, P.Y., Deng, Q., Zakeri, H., Taylor, S.L. and Nickerson, D.A. (1996), 'Increasing the information content of STS-based genome maps: Identifying polymorphisms in mapped STSs', *Genomics* Vol. 31, pp. 123–126.
45. Wang, D.G., Fan, J.B., Siao, C.J., Berno, A. *et al.* (1998), 'Large-scale identification, mapping, and genotyping of single-nucleotide polymorphisms in the human genome', *Science* Vol. 280, pp. 1077–1082.
46. Mattick, J.S. and Makunin, I.V. (2006), 'Non-coding RNA', *Hum. Mol. Genet.* Vol. 15, pp. R17–R29.